



CONTEMPORARY PATHOGENETIC MECHANISM OF SYSTEMIC SCLERODERMA

Akhmedov Khalmurad Sadullaevich, Abdurakhmanova Nargiza Mirza-Bakhtiyarkhanovna, Botirbekov Abdurashid Nazarkulovich
Tashkent medical academy

Article history:	Abstract:
Received: April 4 th 2023 Accepted: May 6 th 2023 Published: June 6 th 2023	Systemic scleroderma is a chronic and often progressive autoimmune inflammation of connective tissue, characterized by fibrosis of the skin and internal organs, common vascular damage with unexplained etiology and heterogeneous clinical manifestations. Systemic scleroderma is a disorder of collagen synthesis and accumulation in the skin (scleroderma) and other internal organs, especially in the connective tissue of the lungs, gastrointestinal tract (gastrointestinal tract), heart and kidneys. Over time, patients show signs of progressive structural and functional impairment of blood vessels and internal organs due to fibrosis. This review article presents current data on the pathogenetic mechanisms of systemic scleroderma.

Keywords: Systemic scleroderma, antinuclear antibodies, T-cells, Raynaud's syndrome, connective tissue growth factor.

INTRODUCTION

Systemic scleroderma is a disease characterized by systemic escalating immune inflammation of the connective tissue and fibro-sclerotic and widespread vasospastic disorders in the skin and subcutaneous tissue, internal organs, i.e. obliterative endarteriolitis [1,10, 28, 29, 28]. Although the concept of scleroderma in general was first described by Lusitanus in 1634 [2,4, 32], the term "scleroderma" was first used by Gintrak in 1847 [5,33]. Although the term "systemic progressive sclerosis" is widely used abroad [7, 22,34], it is now widely accepted as "systemic scleroderma". For many years, scleroderma was considered only a skin disease, and the changes in the internal organs that determine the end of the main disease were overlooked by doctors. Matsui in 1924 [31], in turn, described a widespread degenerative-sclerotic process in connective tissue associated with obliterative disorders in arterioles. The phenomenon shown by the French neurologist Maurice Raynaud in 1862 in a patient and later named Raynaud's disease was actually a symptom associated with vasospastic changes occurring in 96% of systemic scleroderma - Raynaud's syndrome, without knowing it [30]. In fact, it has been confirmed that "Raynaud's syndrome" in systemic scleroderma is one of the main symptoms of the disease.

In the modern world view of this disease, scientists confirm that the main causes of it, as well as a complex process underlie its pathogenesis [9,11,16]. Currently, the causes of systemic scleroderma, that is, the etiology, are unclear. But according to some scientists [10, 29, 19, 28], both endogenous and exogenous factors play a role in the origin of this disease. According to these theories, toxins cause

immune responses by damaging the vascular endothelium, leading to vascular inflammation and tissue sclerosis. In fact, according to studies cited in the literature, fibrosis occurs in the tissues of patients with prolonged contact with polyvinyl chloride [16]. According to the proponents of the influence of infectious factors [13, 17, 18, 28, 24], the immune mechanism develops in the body due to the virus, that is, the immune complexes formed due to antibodies, antigens and complements, without elimination, settle in susceptible tissues and lead to immune inflammation. According to information in the literature [7, 12, 22], factors such as vibration in their profession, long contact with chemicals, dust and long exposure to cold also play an important role in patients suffering from systemic scleroderma. But another group of scientists emphasizes the importance of genetic predisposition in the origin of systemic scleroderma [20, 12].

MODERN VIEWS ON PATHOGENESIS OF SYSTEMIC SCLERODERMA

According to data [14, 16, 23, 35], Ag HLA-10, -V35, -Cw4 association is observed in the genes of patients with systemic scleroderma. In the research of another group of scientists who studied the genetic aspects of the problem [15, 27, 18], 90% of patients with systemic scleroderma show changes in chromosomes: chromatid breaks, edge and circle fragments of chromosomes. In conclusion, diffuse sclerosis occurs in connective tissues as a result of pathological reactions in the immune system under the influence of certain exogenous and endogenous factors in individuals with a genetic predisposition.



It is known that the main three processes in the pathogenesis of systemic scleroderma: metabolism, immune disorders and vascular changes attract the attention of scientists. Given this, what processes occur in metabolism? Metabolic disorder is mainly observed in collagen and its components [32, 17, 29, 24]. Fibrosis is the central process in this joint, and the increased activity of fibroblasts and other collagen-forming cells (including smooth muscle cells in the vessel wall) in the synthesis of collagen, fibronectin, biopolymers - proteoglycan and glycoprotein [16, 32, 13, 19, 23]. According to other available literature sources [26, 28, 25], there is an organic connection between fibroblasts and immunocomplex cells, that is, lymphokines and monokines are involved in controlling the functional activity of fibroblasts. Therefore, in the pathogenesis of systemic scleroderma, the increase in the functional activity of fibroblasts and other collagen-forming cells with a non-stop phenotypic character takes a leading place, which leads to the exacerbation of the pathological process. Also, the role of the autoimmune mechanism in the development of such a fibrosis process is large, that is, the increase in the synthesis of incompletely formed collagens due to the increase in the functional activity of fibroblasts is foreign to the body [12, 18, 16].

This, in turn, causes the appearance of "collagen" autoantigens. According to the information in the literature [21, 13, 14, 25, 18], autoantibodies appear at the expense of those autoantigens, which lead to the renewal of immune complexes. According to the evidence of the established medicine, the detection of antinuclear factor in 90% of patients under observation indicates the involvement of the autoimmune process in the pathogenesis of systemic scleroderma [17, 31, 25]. In addition, T-cells of immunity in the blood: the total amount of T-lymphocytes and their subpopulations - T-suppressor, T-helper decrease, found their place in scientific research of scientists [1, 28, 1, 28, 36]. The increase of circulating immune complexes and autoantibodies in the blood is mainly the result of T-cell deficiency of immunity [18, 26, 28, 19]. Because the lack of T-cells plays a key role in the development of the autoimmune process.

Changes in blood vessels are one of the directions of great importance in the pathogenetic process of systemic scleroderma. As mentioned above, increased functional activity of fibroblasts leads to a specific pathological process in the vessel (endoarteriolitis): microcirculation disorder - changes in serum and cell properties in the blood. According to literature reports [17, 19, 29 37], the cytotoxic effect of

blood increases in this process. As a result, blood solubility increases, collagen synthesis increases, and damage to endothelial cells increases. Therefore, their place is filled by collagen-synthesizing smooth muscle cells, vascular wall hyperplasia and spasms occur [3, 8, 18]. In this process, damage to the vascular endothelium leads to adhesion and aggregation of elements in the blood - leukocytes, erythrocytes, and platelets, that is, blood stasis, intravascular coagulation, and microthrombosis processes take their place [14, 16, 28, 19, 31]. These, in turn, are represented by Raynaud's phenomenon in the clinic of systemic scleroderma [13, 17, 28]. Other scientists who have studied this problem have stated [32, 25, 16, 25] that disturbances in microcirculation flow are the result of changes in immunogenesis. According to the evidence of these scientific works, the formed immune complexes damage the humoral joint of the immune system, especially the vascular endothelium of the compliment system, in the process of settling (fixation) in prone areas. The interaction of immune complexes with the special receptors of platelets in the blood leads to the activation of platelet aggregation properties and blood clotting [4, 7, 14].

In this case, there is the release of biologically active substances into the blood, its swelling and thickening due to plasmatic absorption into the vessel wall, and the deposition of fibrin [13, 6, 18]. Therefore, intravascular narrowing, microcirculation disturbance, platelet adhesion and damage to the vessel wall, in turn, stimulate the migration of "substitute" factors: smooth muscle cells to the vascular intima [7, 27, 19]. In this case, as mentioned above, due to the proliferation of cells, fibroblasts' ability to synthesize collagen increases, and too much secreted collagen leads to perivascular fibrosis [32]. During the phagocytosis of immune complexes, the adhesive property of neutrophils increases, that is, they have a greater chance of sticking to the vessel wall, continuing the inflammatory process [30, 5, 16]. Degranulation of neutrophils during phagocytosis of immune complexes causes the release of lysosomal enzymes. As a result, the basement membrane is damaged, and its remnants turn into autoantigens. This condition ensures the re-emergence of autoantibodies and immune complexes [26].

CONCLUSION

The pathogenesis of systemic scleroderma includes three main mechanisms: fibrosis, autoimmune process, and angiopathy, due to the functional activation of fibroblasts. But in recent years, scientists have been interested in the interrelationship of the



mentioned three processes, the problem of their sequence. It is known that in systemic scleroderma, the main clinical manifestation is the continuous synthesis of low-quality collagens by fibroblasts, which leads to the escalation of the sclerotic process in the skin and internal organs. But collagens themselves also have antigenic properties, and this, in turn, is a driving factor for the development of the autoimmune process. Also, vascular changes are associated with specific disorders of both immune and smooth muscle cells. The pathological changes that we have considered can be thought of as a "chain ring" process. However, the important processes mentioned above do not answer all the questions you want. The reason is that autoimmune diseases are more common in women. According to data [11, 8], autoimmune thyroiditis is 40 times more common in women compared to men, systemic lupus erythematosus is 9 times more common, rheumatoid arthritis and Sjögren's disease are 3 times more common. In particular, women suffer from systemic scleroderma 3 times more than men [31], according to some data [3], 6 times more often. This situation, in turn, attracts attention to gender differences in the pathogenesis of the disease.

Connective tissue growth factor (CTGF), also known as SSN2, is a small secreted protein of the SSN family of three original members, including cysteine-rich (Syr61/ SSN 1), STGF / SSN2, overexpressed in nephroblastoma /SSN 3) STGF is a cysteine-rich extracellular matrix protein composed of four domains or modules. This protein, like other members of the SSN family, contains four different structural modules: an amino terminal insulin-like domain that binds growth factor; cysteine-rich domain; and the carboxyl domain of the terminal cystine node stimulates the synthesis of STGF, a growth factor-like profibrotic cytokine discovered in 1991. Connective tissue growth factor regulates various cellular functions, including the proliferation, migration, adhesion, differentiation and synthesis of extracellular matrix proteins in various types of cells, as well as participates in angiogenesis, chondrogenesis, osteogenesis, wound healing, fibrosis and other more complex biological processes. . An increase in STGF is primarily observed in pathological conditions associated with fibrosis. As an extracellular matrix protein, connective tissue growth factor is thought to integrate various extracellular signals into complex biological responses. The importance of STGF in various fibrosis-related diseases is being actively studied by world scientists, including the fact that this protein plays an important role in the process of fibrosis in lung and heart diseases. However, studying the role

of this protein in the process of skin fibrosis in systemic scleroderma is one of the most urgent issues today.

The main morphological unit of systemic scleroderma is the uncontrolled synthesis of collagens by fibroblasts, resulting in fibrosis in the skin, blood vessels, and internal organs. This, in turn, is related to immunogenesis disorder, and the autoimmune process is its basis.

1. Ахмедов Х. С., Разакова Ф. С., Ахмедова М. С. Значение оптимизации вторичной профилактики системной склеродермии в поддержании качества жизни женщин репродуктивного возраста //Формы и методы социальной работы в различных сферах жизнедеятельности. – 2017. – С. 26-27
2. Ахмедов Х. С. Значение коррекции прогестероновой недостаточности в усилении вторичной профилактики больных системной склеродермией //Врач-аспирант. – 2007. – Т. 17. – №. 2. – С. 92-96.
3. Ботирбеков А.Н., Ахмедов Х.С., Абдурахманова Н.М. Тизимли склеродермиянинг замонавий патогенетик механизмлари// Инфекция, иммунитет ва фармакология. -2023
4. Зарипов С.И., Ахмедов Х.С., Абдурахманова Н.М. Значение аутоантел в патогенезе системной склеродермии. -2023
5. Рихсиева Л. М., Абдурахманова Н. М. Б., Ахмедов Х. С. Оценка уровня интерлейкина-17 у больных аксиальным спондилоартритом. – 2023.
6. Соболевский П., Маслинская М., Вечорек М., Лагун З., Малевска А., Рашкевич М., Ницкович Р., Шиманская Э. и Валецка И. (2019). Системный склероз - мультидисциплинарное заболевание: клиника и лечение. Ревматология, 57 (4), 221–233.
7. Akhmedov, K. S., Rixsiyeva, L. M., Abdurahmanova, N. M., Botirbekov, A. N., Turaev, I. A., & Raximov, S. S. (2023). Changes in C-TELOPEPTIDE COLLAGEN-II level in patients with ankylosing spondyloarthritis after COVID-19. Journal of Modern Educational Achievements, 3(3), 78-87.
8. Axmedov Kh.S, Abdurakhmanova N.M, Zaripov S.S., & Turaev I.A. (2023). THE EFFECT OF CLIMATE-GEOGRAPHICAL FACTORS ON RHEUMATOID ARTHRITIS ACTIVITY. World



- Bulletin of Public Health, 18, 67-69. Retrieved from
9. Abdurakhmanova N., Akhmedov Kh. Effect of pro-inflammatory cytokine-Interleukin- 6 on the course of ankylosing spondylitis in patients after COVID-19 //Annals of the Rheumatic Diseases. – 2022. – C. 1533-1533.
 10. Abdurakhmanova N. M. et al. Modern methods of treatment of patients with ankylosing spondylitis //International Journal of Advance Scientific Research. – 2022. – T. 2. – №. 11. – C. 112-118
 11. Abdurakhmanova N. M. High concentration of tumor necrosis factor in ankylosing spondylitis patients after COVID-19 //British medical journal. – 2022. – T. 2. – №. 1.2.
 12. Akhmedov K., Abdurakhmanova N., Buranova S. Features of the clinical course of rheumatoid spine against the background of the influence of xenobiotics //American Journal of Interdisciplinary Research and Development. – 2023. – T. 12. – C. 142-147.
 13. Abdurakhmanova, N. M., & Ahmedov, K. S. (2022). Influence of accepting basic antirheumatic therapy for ankylosing spondyloarthritis on the clinical course Covid-19. *ACADEMICIA: An International Multidisciplinary Research Journal*, 12(7), 71-76.
 14. Barsotti S, Bellando Randone S, Guiducci S, et al. Systemic sclerosis: a critical digest of the recent literature. *Clin Exp Rheumatol* 2014; 32: S194-S205.
 15. Bertrand D, Dehay J, Ott J, et al. Kidney transplantation in patients with systemic sclerosis: a nationwide multicentre study. *Transpl Int* 2017; 30: 256-265
 16. Castro SV, Jimenez SA. Biomarkers in systemic sclerosis. *Biomark Med* 2010; 4: 133-147. / Barsotii S, Stagnaro C, Della Rossa A. Systemic sclerosis: a critical digest of the recent literature. *Clin Exp Rheumatol* 2105; 33 (Suppl 91): S3-14.
 17. Cavazzana, I., Vojinovic, T., Airo', P. et al. Systemic Sclerosis-Specific Antibodies: Novel and Classical Biomarkers. *Clinic Rev Allerg Immunol* (2022).
 18. Clark, K.E.N., Campochiaro, C., Host, L.V. et al. Combinations of scleroderma hallmark autoantibodies associate with distinct clinical phenotypes. *Sci Rep* 12, 11212 (2022)
 19. Dieudé P, Guedj M, Wipff J, et al. BANK1 is a genetic risk factor for diffuse cutaneous systemic sclerosis and has additive effects with IRF5 and STAT4. *Arthritis Rheum* 2009; 60: 3447-3454.
 20. Emmanuel A. Current management of the gastrointestinal complications of systemic sclerosis. *Nat Rev Gastroenterol Hepatol* 2016; 13: 461-472
 21. Furue M, Mitoma C, Mitoma H, et al. Pathogenesis of systemic sclerosis – current concept and emerging treatments. *Immunol Res* 2017; 65: 790-797.
 22. Gong, P., Ding, Y., Sun, R. et al. Mesenchymal stem cells alleviate systemic sclerosis by inhibiting the recruitment of pathogenic macrophages. *Cell Death Discov.* 8, 466 (2022)
 23. Hasegawa M. Biomarkers in systemic sclerosis: their potential to predict clinical courses. *J Dermatol* 2016; 43: 29-38.
 24. HOFFMANN-VOLD AM, ALLANORE Y, ALVESM et al.: Progressive interstitial lung disease in patients with systemic sclerosis-associated interstitial lung disease in the EUSTAR database. *Ann Rheum Dis* 2021; 80: 219-27
 25. Jerjen R, Nikpour M, Krieg T, Denton CP, Saracino AM. Systemic sclerosis in adults. Part I: Clinical features and pathogenesis. *J Am Acad Dermatol.* 2022 Nov;87(5):937-954. doi: 10.1016/j.jaad.2021.10.065. Epub 2022 Feb 4. PMID: 35131402.
 26. Jin J, Chou C, Lima M, et al. Systemic Sclerosis is a Complex Disease Associated Mainly with Immune Regulatory and Inflammatory Genes. *Open Rheumatol J* 2014; 8: 29-42.
 27. Kayser C, Fritzler MJ. Autoantibodies in systemic sclerosis: unanswered questions. *Front Immunol.* 2015 Apr 15;6:167. doi: 10.3389/fimmu.2015.00167. PMID: 25926833; PMCID: PMC4397862.
 28. Keppeke GD, Satoh M, Kayser C, Matos P, Hasegawa T, Tanaka S, Diogenes L, Amaral RQ, Rodrigues SH and Andrade LEC (2022) A cell-based assay for detection of anti-fibrillarin autoantibodies with performance equivalent to immunoprecipitation. *Front. Immunol.* 13:1011110. doi: 10.3389/fimmu.2022.1011110
 29. Kim S, Park HJ, Lee SI. The Microbiome in Systemic Sclerosis: Pathophysiology and Therapeutic Potential. *Int J Mol Sci.* 2022 Dec 18;23(24):16154. doi:



- 10.3390/ijms232416154. PMID: 36555792; PMCID: PMC9853331.
30. Kowal-Bielecka O, Kuryliszyn-Moskal A. Twardzina układowa. *Reumatologia* 2016; Suppl 1: 51-55.
 31. Kucharz EJ, Kopeć-Mędrek M. Systemic sclerosis sine scleroderma. *Adv Clin Exp Med* 2017; 26: 875-880.
 32. Lee A. Fibrillarin antibodies in systemic sclerosis. *Arch Rheumatol.* 2021 Oct 18;37(2):311-312. doi: 10.46497/ArchRheumatol.2022.9038. PMID: 36017207; PMCID: PMC9377177.
 33. Mejia Otero C, Assassi S, Hudson M, Mayes MD, Estrada-Y-Martin R, Pedroza C, Mills TW, Walker J, Baron M, Stevens W, Proudman SM, Nikpour M, Mehra S, Wang M, Fritzler MJ; Canadian Scleroderma Research Group; Australian Scleroderma Cohort Study; Genetics versus Environment in Scleroderma Outcome Study. Antifibrillarin Antibodies Are Associated with Native North American Ethnicity and Poorer Survival in Systemic Sclerosis. *J Rheumatol.* 2017 Jun;44(6):799-805. doi: 10.3899/jrheum.160574. Epub 2017 Apr 1. PMID: 28365584; PMCID: PMC5457664.
 34. Mirza-Bakhtiyarxonovna A. N., Sadullaevich A. K., Samandarovich R. S. impact of COVID-19 on the course of ankylosing spondylitis //Journal of Modern Educational Achievements. – 2023. – T. 6. – №. 6. – C. 202-207.
 35. Schiopu E, Impens AJ, Phillips K. Digital ischemia in scleroderma spectrum of diseases. *Int J Rheumatol* 2010; 2010: 923743.
 36. Steen VD, Medsger TA. Severe organ involvement in systemic sclerosis with diffuse scleroderma. *Arthritis Rheum* 2000; 43:2437-2444.
 37. TERRIER B, DECHARTRES A, GOUYA H et al.: Cardiac intravoxel incoherent motion diffusion-weighted magnetic resonance imaging with T1 mapping to assess myocardial perfusion and fibrosis in systemic sclerosis: association with cardiac events from a prospective cohort study. *Arthritis Rheumatol* 2020; 72: 1571-80.
 38. Van den Hoogen F, Khanna D, Fransen J, et al. 2013 Classification Criteria for Systemic Sclerosis: an American College of Rheumatology/ European League Against Rheumatism collaborative initiative. *Ann Rheum Dis* 2013; 72: 1747-755.