

Available Online at: https://www.scholarexpress.net

Volume-38, September 2024

ISSN: 2749-3644

# PREDICTION OF HIV PREVALENCE AMONG INDIVIDUALS AGED 15-49 YEARS IN RWANDA USING HOLT'S LINEAR METHOD

### Dr. Smartson. P. NYONI¹, Thabani NYONI²

<sup>1</sup>ZICHIRe Project, University of Zimbabwe, Harare, Zimbabwe <sup>2</sup>Independent Researcher & Health Economist, Harare, Zimbabwe

Article history: Abstract:	
Received: June 20 <sup>th</sup> 2024 Accepted: July 14 <sup>h</sup> 2024	This study uses annual time series data of HIV prevalence among individuals aged 15-49 years for Rwanda from 1990 to 2020 to predict future trends of HIV prevalence over the period 2021 to 2030. The study utilizes Holt's linear exponential smoothing model. The optimal values of smoothing constants a and $\beta$ are 0.9 and 0.1 respectively based on minimum MSE. The results of the study indicate that annual HIV prevalence among individuals aged 15-49 years will continue to decline over the out of sample period. Therefore, we encourage authorities to address main drivers of HIV transmission among this age group. There is need to improve HIV case detection and prevention among high risk groups

**Keywords:** Exponential smoothing, Forecasting, HIV prevalence

#### **BACKGROUND**

HIV remains a public health concern globally, causing negative health, social and economic consequences among all age groups in low-middle income countries (Kawuki et al. 2022; Negesse et al. 2021). According to UNAIDS data, around 38 million people were living with HIV (PLWH) worldwide in 2022. Africa reported the highest prevalence of HIV with an estimated average of 3.9% of its population living with HIV (Mbabazi et al. 2022; Operto, 2020). Young women and teenage girls account for 25% of new infections in Sub-Saharan Africa (SSA) (UNAIDS, 2020). Eastern and Southern African women face the highest burden of HIV contributing to three in five new infections, while young women aged 15 to 24 years are almost three times more likely than their male peers to acquire HIV infection (UNAIDS, 2020; Niragire et al. 2021). According to UNICEF, approximately more than 700 adolescents acquire HIV daily and over 360,000 are expected to die of HIV/AIDS related illnesses between 2018 and 2030. Adolescent HIV/AIDS-related mortality in Africa remains significant and AIDS is among the leading causes of death among African adolescents with SSA having the highest AIDS related mortality burden (Badru et al. 2017; UNICEF, 2015; WHO, 2014). Rwanda reported an HIV prevalence of 3.0% in 2019 among people aged 15-49 years and had a higher proportion of HIV in women (3.6%) than in men (2.2%) (Niragire et al. 2015). HIV prevalence in Rwanda is still

low compared to other East African countries, however HIV risk factors continue to pose a huge threat to the upsurge of new HIV infections (Broderick et al. 2021; Niragire et al. 2021). Factors which predict HIV infection include sexual behaviors such as concurrent sexual partners, an earlier sex debut (< 19 years), and noncondom use. Furthermore, women with a history of sexually transmitted infections (STIs) living in woman headed households were more susceptible to HIV infection (Negesse et al. 2021). Limited comprehensive knowledge of HIV has been reported as one of the major factors associated with high prevalence of HIV among adolescents and young women (AGYW) (Frimpong et al. 2021; Siziya et al. 2008). Having adequate comprehensive HIV knowledge has been found to enhance adolescents' and young women's ability to negotiate for safer sex hence helping to reduce the risk of contracting HIV (Frimpong et al. 2021; De Coninck et al. 2014).

The objective of this paper is to model and forecast HIV prevalence among individuals aged 15-49 years for Rwanda using Holt's linear exponential smoothing technique. The findings of this research are anticipated to guide policy, planning and allocation of resources towards targeted HIV prevention, treatment and support programs in Rwanda with the aim of curbing new HIV infections and reducing HIV/AIDS related morbidity and mortality

.



World Bulletin of Public Health (WBPH)
Available Online at: https://www.scholarexpress.net
Volume-38, September 2024
ISSN: 2749-3644

# **LITERATURE REVIEW**

Author(s)	Objective (s)	Methodology	Key finding (s)
Niyompano et al. (2024)	To determine the prevalence of never being tested for HIV and its associated factors among sexually active individuals aged 15–56 who participated in the Rwanda AIDS Indicators and HIV Incidence Survey (RAIHIS).	-cross-sectional study -Applied Bivariate and multivariate logistic regression models	The prevalence of non-testing for HIV was 17.37%. Being aged 15–30 years (aOR 2.57, 95%CI 1.49–4.43, p < 0.001) and male (aOR 2.44, 95%CI 1.77–3.36, p < 0.001) was associated with an increase in the odds of never testing for HIV
Kawuki et al. (2023)	To determine the prevalence of the risk factors for the acquisition of HIV among sexually active women in Rwanda.	- Secondary data from the 2020 Rwanda Demographic Health Survey, comprising 10,684 sexually active women, was used -Multivariable logistic regression was conducted to determine the associated risk factors using the SPSS (version 25).	More than a quarter of sexually active women in Rwanda had exposure to at least one risk factor for HIV acquisition.
Kawuki et al. (2023)	To assess the prevalence of comprehensive knowledge about HIV/AIDS and associated factors among adolescent girls in Rwanda	-used secondary data from the Rwanda Demographic and Health Survey (RDHS) 2020 comprising 3258 adolescent girls (aged 15 to 19 years)Conducted multivariable logistic regression to explore the associated factors, using SPSS (version 25).	Adolescent girls with secondary education (AOR = 1.40, 95% CI: 1.13–3.20), health insurance (AOR = 1.39, 95% CI: 1.12–1.73), a mobile phone (AOR = 1.26, 95% CI: 1.04–1.52), exposure to television (AOR = 1.23, 95% CI: 1.05–1.44), and a history of an HIV test (AOR = 1.26, 95% CI: 1.07–1.49) had higher odds of comprehensive HIV knowledge, compared to their respective counterparts
Nsanzimana et al. (2022)	To measure national HIV Incidence and prevalence. District- level Estimates were modeled to inform resources allocation	The 2018–2019 Rwanda Population– based HIV Impact Assessment (RPHIA) was conducted	HIV Prevalence among adults in Rwanda Is 3.0%; HIV Incidence is low at 0.08%. District-level Modeling has identified disproportionately affected urban



Available Online at: https://www.scholarexpress.net

Volume-38, September 2024

ISSN: 2749-3644

			hotspots: areas to focus resources.
Biraguma et al. (2018)	To determine the association between physical and mental health-related dimensions of quality of life (QOL) with behavioral and biological risk factors, after controlling sociodemographic and HIV-related factors in adults living with HIV in Rwanda.	-Cross-sectional study -using the WHO STEPwise approach and Kinyarwanda version of the MOS-HIV Health Survey, risk factors for NCDs and HRQOL were analyzed for 794 PLWH, both HIV+ on ART and ART-naïve. Multiple regression analysis was used to examine the relationship between CMD risk factors and physical health and mental health summary scores.	Behavioral and biological risk factors for NCDs were significantly associated with a lower HRQOL.
Nsanzimana et al. (2017)	to characterize HIV incidence across Rwanda	prospective HIV incidence survey for the period of 2013–14	The incidence of HIV in Rwanda was higher than that previously estimated from models, with outbreaks seeming to contribute to the ongoing epidemic
Nsanzimana et al. (2015)	to assess the effect of increased access to antiretroviral therapy on life expectancy among HIV-positive patients in two distinct periods of lower and higher antiretroviral therapy coverage (1997–2007 and 2008–11).	retrospective observational cohort study	Life expectancy at 20 years of age in the period of 1997–2007 was 20·4 additional years (95% CI 19·5–21·3); for the period of 2008–11, life expectancy had increased to 25·6 additional years (95% CI 24·8–26·4).

#### **METHODOLOGY**

This study utilizes an exponential smoothing technique to model and forecast future trends of HIV prevalence among individuals aged 15-49 years in Rwanda. In exponential smoothing forecasts are generated from the smoothed original series with the most recent historical values having more influence than those in the more distant past as more recent values are allocated more weights than those in the distant past. This study uses the Holt's linear method (Double exponential smoothing) because it is an appropriate technique for modeling linear data.

Holt's linear method is specified as follows:

Model equation

 $A_t = \mu_t + \rho_t \mathbf{t} + \varepsilon_t$ <u>Smoothing equation</u>

 $S_t = \alpha A_t + (1-\alpha) (S_{t-1} + b_{t-1})$ 

0<∝<1

Trend estimation equation

 $b_t = \beta (S_t - S_{t-1}) + (1 - \beta)b_{t-1}$ 0<\beta<1

0<*p*<1

Forecasting equation

 $f_{t+h} = S_t + hb_t$ 

 $A_t$  is the actual value of HIV prevalence at time t

 $\varepsilon_t$  is the time varying **error term** 

 $\mu_t$  is the time varying mean (**level**) term

 $\rho_t$  is the time varying **slope term** 

**t** is the trend component of the time series



Available Online at: https://www.scholarexpress.net

Volume-38, September 2024

ISSN: 2749-3644

 $\mathcal{S}_t$  is the exponentially smoothed value of HIV prevalence at time t

 $\alpha$  is the exponential smoothing constant for the data  $\beta$  is the smoothing constant for trend  $f_{t+h}$  is the h step ahead forecast

 $b_t$  is the trend estimate (slope of the trend) at time t  $b_{t-1}$  is the trend estimate at time t-1

**Data Issues** 

Table 1: ES model summary

This study is based on annual HIV prevalence among individuals aged 15-49 years in Rwanda for the period 1990 – 2020. The out-of-sample forecast covers the period 2021 – 2030. All the data employed in this research paper was gathered from the World Bank online database.

#### **Findings of the study**

**Exponential smoothing Model Summary** 

Variable	A
Included Observations	31
Smoothing constants	
Alpha (a) for data	0.900
Beta (β) for trend	0.100
Forecast performance measures	
Mean Absolute Error (MAE)	0.207329
Sum Square Error (SSE)	4.485410
Mean Square Error (MSE)	0.144691
Mean Percentage Error (MPE)	-1.119141
Mean Absolute Percentage Error (MAPE)	6.930046

Residual Analysis for the Applied Model




Figure 1: Residual analysis

In-sample Forecast for A



Available Online at: https://www.scholarexpress.net

Volume-38, September 2024

ISSN: 2749-3644




Figure 2: In-sample forecast for the A series

# Actual and Smoothed graph for A series

Figure 3: Actual and smoothed graph for A series **Actual and Smoothed Graph** 5.0 4.5 4.0 3.5 3.0 2.5 2.0 1.5 1990 1993 1996 1999 2002 2005 2008 2011 2014 2017 2020 Time

Out-of-Sample Forecast for A: Actual and Forecasted Graph


Figure 4: Out-of-sample forecast for A: actual and forecasted graph



Available Online at: https://www.scholarexpress.net

Volume-38, September 2024

ISSN: 2749-3644



Out-of-Sample Forecast for A: Forecasts only Table 2: Tabulated out-of-sample forecasts

Year	Forecasted HIV prevalence
2021	2.4353
2022	2.3669
2023	2.2984
2024	2.2300
2025	2.1615
2026	2.0930
2027	2.0246
2028	1.9561
2029	1.8877
2030	1.8192

The main results of the study are shown in table 1. It is clear that the model is stable as confirmed by evaluation criterion as well as the residual plot of the model shown in figure 1. It is projected that annual HIV prevalence among individuals aged 15-49 years will continue to decline over the out of sample period.

#### **Policy implication and conclusion**

Our model projections indicate that annual HIV prevalence among individuals aged 15-49 years will continue to decline over the out of sample period. Therefore, this research calls for the authorities to address main drivers of HIV transmission among this age group. There is need to improve HIV case detection and prevention among high risk groups.

#### **REFERENCES**

[1] Kawuki J, Kamara K & Sserwanja Q (2022). Prevalence of risk factors for human immunodeficiency virus among women of reproductive age in Sierra Leone: a 2019 nationwide survey. BMC Infect Dis. 22(1):1–9.

- [2] Negesse Y, Mankelkl G, Setegn M & Fetene G (2021). Multilevel analysis of factors associated with HIV among women of reproductive age (15–49 years old) in Ethiopia: bayesian approach. Women's Health. 17:17455065211067638.
- [3] UNAIDS (2022). In Danger: UNAIDS Global AIDS Update 2022 2022 [Available from: https://www.unaids.org/en/resources/docume nts/2022/ in-danger-global-aids-update
- [4] Operto E (2020). Knowledge, attitudes, and practices regarding exclusive breastfeeding among HIV-positive mothers in Uganda: a qualitative study. Int J Health Plann Manag. 35(4):888–96.
- [5] Mbabazi L, Nabaggala MS, Kiwanuka S, Kiguli J, Laker E & Kiconco A (2022). Factors associated with uptake of contraceptives among HIV positive women ondolutegravir based antiretroviral treatment-a cross sectional survey in



Available Online at: https://www.scholarexpress.net

Volume-38, September 2024

ISSN: 2749-3644

- urban Uganda. BMC Womens Health. 22(1):1–8.
- [6] Niragire F, Achia TN, Lyambabaje A & Ntaganira J (2015). Bayesian mapping of HIV Infection among women of reproductive age in Rwanda. PLoS ONE. 10(3):e0119944.
- [7] Niragire F, Ndikumana C, Nyirahabimana MG, Uwizeye D (2021). Prevalence and factors associated with fertility desire among HIV-positive women in Rwanda in the context of improved life expectancy. Archives of Public Health. 79(1):1–10.
- [8] Broderick K, Ponticiello M, Nabukalu D, Tushemereirwe P, Nuwagaba G, & King R (2021). Shortening the road to improve engagement with HIV testing resources: a qualitative study among stakeholders in rural Uganda. AIDS Patient Care STDs. 35(2):56–62.
- [9] Joint United Nations Programme on HIV/AIDS (UNAIDS). UNAIDS 2020 | Reference: UNAIDS Data 2020- State of the Epidemic. Geneva: UNAIDS. ; 2020. Available from: <a href="https://www.unaids.org/sites/default/files/med">https://www.unaids.org/sites/default/files/med</a> ia asset/2020 aids-data-book en.pdf
- [10] UNICEF. Around 80 adolescents will die of AIDS every day by 2030, at current trends UNICEF. <a href="https://www.unicef.org/press-releases/around-80-adolescents-will-die-aids-every-day-2030-current-trends-unicef">https://www.unicef.org/press-releases/around-80-adolescents-will-die-aids-every-day-2030-current-trends-unicef</a>.
- [11] Badru T, Mwaisaka J, Khamofu H, Agbakwuru C, Adedokun O, Pandey SR, Essiet P, James E, Chen-Carrington A, Mastro TD, et al. HIV comprehensive knowledge and prevalence among young adolescents in Nigeria: evidence from Akwa Ibom AIDS indicator survey, 2017. BMC Public Health. 2020;20(1):45.
- [12] UNICEF, Adolescent. deaths from AIDS tripled since 2000. Press release. 2015. https://www.efe.com/efe/english/life/unicef-says-adolescent-deaths-fromaids-tripled-since-2000/50000263-2775480.
- [13] World Health Organization. Health for the world's adolescents: a second chance in the second decade: summary. World Health Organization. ; 2014. https://apps.who.int/iris/handle/10665/112750
- [14] Frimpong JB, Budu E, Adu C, Mohammed A, Tetteh JK, Seidu A-A & Ahinkorah BO (2021). Comprehensive HIV/AIDS knowledge and safer sex negotiation

- among adolescent girls and young women in sub-Saharan Africa. J Biosoc Sci 2021:1–13.
- [15] Siziya S, Muula AS & Rudatsikira E (2008). HIV and AIDS-related knowledge among women in Iraq. BMC Res Notes. 1(1):123
- [16] De Coninck Z, Feyissa IA, Ekström AM & Marrone G (2014). Improved HIV awareness and perceived empowerment to Negotiate Safe Sex among Married Women in Ethiopia between 2005 and 2011. PLoS ONE. 9(12):e115453